

November 1991, DSF-91/20

α -SURFACES FOR COMPLEX SPACE-TIMES
WITH TORSION

Giampiero Esposito

Istituto Nazionale di Fisica Nucleare, Sezione di Napoli, Gruppo IV,

Mostra d'Oltremare, Padiglione 20, 80125 Napoli

Dipartimento di Scienze Fisiche,

Mostra d'Oltremare, Padiglione 19, 80125 Napoli

Abstract. - This paper studies necessary conditions for the existence of α -surfaces in complex space-time manifolds with nonvanishing torsion. For these manifolds, Lie brackets of vector fields and spinor Ricci identities contain explicitly the effects of torsion. This leads to an integrability condition for α -surfaces which does not involve just the self-dual Weyl spinor, as in complexified general relativity, but also the torsion spinor, in a nonlinear way, and its covariant derivative. Interestingly, a particular solution of the integrability condition is given by right-flat and right-torsion-free space-times.

PACS 04.20.Cv - Fundamental problems and general formalism.

PACS 04.50 - Unified field theories and other theories of gravitation.

α -Surfaces for Complex Space-Times with Torsion

Twistor theory was created by Roger Penrose as an approach to the quantum-gravity problem where null surfaces and some particular complex manifolds are regarded as fundamental entities, whereas space-time points are only derived objects, since they might become ill-defined at the Planck length [1]. It is by now well-known that the building blocks of classical field theory in Minkowski space-time are the α -planes. In other words, one first takes complexified compactified Minkowski space $CM^\#$, and one then defines α -planes as null two-surfaces, such that the metric vanishes over them, and their null tangent vectors have the two-component-spinor form $\lambda^A \pi^{A'}$, where λ^A is varying and $\pi^{A'}$ is fixed by a well-known differential equation [1]. This definition can be generalized to complex or real Riemannian (i.e. with positive-definite metric) space-times provided the Weyl curvature is anti-self-dual, giving rise to the so-called α -surfaces [1].

We are here interested in the case of complex space-times with nonvanishing torsion (hereafter referred to as CU_4 space-times). This study appears relevant at least for the following reasons: (i) the definition of torsion is a peculiarity of relativistic theories of gravitation; (ii) the gauge theory of the Poincaré group leads to theories with torsion; (iii) theories with torsion are theories of gravity with second-class constraints; (iv) if torsion is nonvanishing, the occurrence of cosmological singularities can be less generic than in general relativity [2]. We have thus studied the problem: what are the conditions on curvature and torsion for a CU_4 space-time to admit α -surfaces (defined as above) ?

The starting point of our calculation, based on the use of the full U_4 -connection with torsion, is the evaluation of the Lie bracket of two vector fields X and Y tangent to a

totally null two-surface in CU_4 . By virtue of Frobenius' theorem [1], this Lie bracket is a linear combination of X and Y

$$[X, Y] = \varphi X + \rho Y \quad , \quad (1)$$

where φ and ρ are scalar functions. Moreover, using the definition of the torsion tensor $S(X, Y)$, one also has

$$[X, Y] \equiv \nabla_X Y - \nabla_Y X - 2S(X, Y) \quad . \quad (2)$$

Thus, since in CU_4 models the torsion tensor, antisymmetric in the first two indices, can be expressed spinorially as

$$S_{ab}{}^c = \chi_{AB}{}^{CC'} \epsilon_{A'B'} + \tilde{\chi}_{A'B'}{}^{CC'} \epsilon_{AB} \quad , \quad (3)$$

where the spinors χ and $\tilde{\chi}$ are symmetric in AB and $A'B'$ respectively, and are *totally independent*, one finds by comparison of Eqs. (1) and (2) that

$$\pi^{A'} \left(\nabla_{AA'} \pi_{B'} \right) = \xi_A \pi_{B'} - 2\pi^{A'} \pi^{C'} \tilde{\chi}_{A'B'AC'} \quad , \quad (4)$$

where we have set $X^a = \lambda^A \pi^{A'}$, $Y^a = \mu^A \pi^{A'}$, while φ and ρ are obtained as follows: $-\mu^A \xi_A = \varphi$, $\lambda^A \xi_A = \rho$. This is the desired necessary condition for the field $\pi^{A'}$ to define an α -surface in the presence of torsion. We now have to derive the integrability condition for Eq. (4). For this purpose, we operate with $\pi^{B'} \pi^{C'} \nabla_{C'}^A$ on both sides of Eq. (4), we

repeatedly use the Leibniz rule and Eq. (4), and we also use the spinor formula for the Riemann tensor and spinor Ricci identities. The relations we need are

$$\begin{aligned}
 R_{abcd} = & \psi_{ABCD}\epsilon_{A'B'}\epsilon_{C'D'} + \tilde{\psi}_{A'B'C'D'}\epsilon_{AB}\epsilon_{CD} \\
 & + \Phi_{ABC'D'}\epsilon_{A'B'}\epsilon_{CD} + \tilde{\Phi}_{A'B'CD}\epsilon_{AB}\epsilon_{C'D'} \\
 & + \Sigma_{AB}\epsilon_{A'B'}\epsilon_{CD}\epsilon_{C'D'} + \tilde{\Sigma}_{A'B'}\epsilon_{AB}\epsilon_{CD}\epsilon_{C'D'} \\
 & + \Lambda\left(\epsilon_{AC}\epsilon_{BD} + \epsilon_{AD}\epsilon_{BC}\right)\epsilon_{A'B'}\epsilon_{C'D'} \\
 & + \tilde{\Lambda}\left(\epsilon_{A'C'}\epsilon_{B'D'} + \epsilon_{A'D'}\epsilon_{B'C'}\right)\epsilon_{AB}\epsilon_{CD} \quad , \tag{5}
 \end{aligned}$$

$$\left[\nabla_{C(A'}\nabla^C_{B')} - 2\tilde{\chi}_{A'B'}{}^{HH'}\nabla_{HH'} \right] \pi^{C'} = \tilde{\psi}_{A'B'E'}{}^{C'} \pi^{E'} - 2\tilde{\Lambda}\pi_{(A'}\epsilon_{B')}^{C'} + \tilde{\Sigma}_{A'B'}\pi^{C'} \quad . \tag{6}$$

As usual, the *twiddle* symbol denotes spinor or scalar quantities independent of their untwiddled counterpart. The spinors ψ and $\tilde{\psi}$ are the anti-self-dual and self-dual Weyl spinors respectively, and are thus invariant under conformal rescalings of the metric. The symmetric spinors Σ and $\tilde{\Sigma}$ express the antisymmetric part of the Ricci tensor: $R_{[ab]} = \Sigma_{AB}\epsilon_{A'B'} + \tilde{\Sigma}_{A'B'}\epsilon_{AB}$. More details on the spinors Φ , $\tilde{\Phi}$ and the scalar functions Λ , $\tilde{\Lambda}$ can be found in Ref. [3]. Eq. (6) is proved defining the operator $\square_{ab} \equiv 2\nabla_{[a}\nabla_{b]} - 2S_{ab}{}^c \nabla_c$ and the self-dual null bivector $k^{ab} \equiv \kappa^A\kappa^B\epsilon^{A'B'}$, and using the tensor form of the Ricci identity for U_4 theories

$$\square_{ab}k^{cd} = R_{abe}{}^c k^{ed} + R_{abe}{}^d k^{ce} \quad . \tag{7}$$

Thus, using the identity $2\nabla_{[a}\nabla_{b]} = \epsilon_{A'B'}\square_{AB} + \epsilon_{AB}\square_{A'B'}$, where $\square_{C'A'} \equiv \nabla_{A(C'}\nabla^A_{A')}$, a lengthy calculation of Eq. (7) yields the spinorial equations

$$\left[\square_{AB} - 2\chi_{AB}{}^{HH'}\nabla_{HH'} \right] \kappa^C = \psi_{ABE}{}^C \kappa^E - 2\Lambda\kappa_{(A}\epsilon_{B)}^C + \Sigma_{AB}\kappa^C \quad , \tag{8}$$

$$\left[\square_{A'B'} - 2\tilde{\chi}_{A'B'}{}^{HH'} \nabla_{HH'} \right] \kappa^C = \tilde{\Phi}_{A'B'E}{}^C \kappa^E + \tilde{\Sigma}_{A'B'} \kappa^C \quad . \quad (9)$$

Eq. (6) is then obtained replacing unprimed indices with primed indices, untwiddled spinors and scalars with twiddled spinors and scalars, and κ^C with $\pi^{C'}$ in Eq. (8).

Finally, using Eq. (4), Eq. (6), the identity

$$\begin{aligned} \pi^{B'} \pi^{C'} \nabla_{C'}^A \left[\pi^{A'} \left(\nabla_{AA'} \pi_{B'} \right) \right] &= \pi^{B'} \pi^{C'} \left(\nabla_{C'}^A \pi^{A'} \right) \left(\nabla_{AA'} \pi_{B'} \right) \\ &\quad - \pi^{A'} \pi^{B'} \pi^{C'} \left(\square_{C'A'} \pi_{B'} \right) \quad , \end{aligned} \quad (10)$$

and the well-known property $\lambda_A \lambda^A = \pi_{A'} \pi^{A'} = 0$, the integrability condition for α -surfaces is found to be

$$\begin{aligned} \tilde{\psi}_{A'B'C'D'} &= -4\tilde{\chi}_{A'B'AL'} \tilde{\chi}_{C'}{}^{L'A}{}_{D'} + 4\tilde{\chi}_{L'B'AC'} \tilde{\chi}_{A'D'}{}^{AL'} \\ &\quad + 2\nabla_{D'}^A \left(\tilde{\chi}_{A'B'AC'} \right) \quad . \end{aligned} \quad (11)$$

Note that an analogous result holds if the metric is positive-definite rather than complex. A four-manifold with nonvanishing torsion and positive-definite metric is here denoted by RU_4 . Interestingly, a particular solution of Eq. (11) is given by $\tilde{\psi}_{A'B'C'D'} = 0$, $\tilde{\chi}_{A'C'AB'} = 0$. By analogy with complexified general relativity, the particular CU_4 and RU_4 space-times satisfying these additional conditions are here called right-flat and right-torsion-free (although our definition does not involve the Ricci tensor, and is therefore different from Eq. (6.2.1) of Ref. [1]). This means that the surviving Weyl and torsion spinors, i.e. ψ_{ABCD} and $\chi_{AB}{}^{CC'}$, do not affect the integrability condition (11) for α -surfaces. Note also that this is only possible for CU_4 and RU_4 models of gravity, since only for these theories the torsion spinors χ and $\tilde{\chi}$, and the Weyl spinors ψ and $\tilde{\psi}$, are totally independent [1].

Acknowledgements

We are much indebted to the referees of the journal *General Relativity and Gravitation* for suggesting the correct forms of the result and correcting numerous other errors in the original version of this paper, and to a referee of *Nuovo Cimento* for helpful comments which improved style and nature of our work. We are also grateful to Professors Giuseppe Marmo and Ruggiero de Ritis for reading the manuscript and many enlightening conversations, and to Dr. David Hughes for correspondence on spinor methods.

References

- [1] R. S. WARD and R. O. WELLS: *Twistor Geometry and Field Theory* (Cambridge University Press, Cambridge, 1990).
- [2] G. ESPOSITO: *Quantum Gravity, Quantum Cosmology and Lorentzian Geometries* (Springer-Verlag, Berlin, 1992).
- [3] R. PENROSE: *Found. Phys.*, **13**, 325 (1983).

RIASSUNTO

Questo lavoro studia condizioni necessarie per l'esistenza di α -superfici in modelli di spazio-tempo complesso con torsione non nulla. Per queste varietà, le parentesi di Lie di campi vettoriali e le identità spinoriali di Ricci contengono esplicitamente gli effetti della torsione. Questo conduce ad una condizione di integrabilità per le α -superfici che non involve soltanto lo spinore di Weyl auto-duale, come nella relatività generale complessificata, ma anche lo spinore di torsione, in modo non lineare, e la sua derivata covariante. Di un certo interesse, una soluzione particolare della condizione di integrabilità è data da modelli di spazio-tempo piatti a destra e liberi da torsione a destra.